

Gluon coupling strength in H_{QCD} with small cutoffs

Stanisław D. Glazek^{a,*}

^a Institute of Theoretical Physics, Warsaw University,
ul. Hoża 69, 00-681 Warsaw, Poland

This contribution presents the running triple-gluon-vertex coupling constant, g_λ , in Hamiltonians for the gluons that are characterized by the size $\sim 1/\lambda$. The coupling constant is obtained from renormalization group equations for effective particles in canonical light-front QCD in third order perturbation theory. λ plays the role of a finite cutoff parameter and is varied from 100 GeV down to 100 MeV.

1. Introduction

A renormalization group approach to light-front (LF) Hamiltonian QCD was outlined by Wilson et al. in Ref. [1] for the purpose of deriving hadronic structure in terms of well defined wave functions for constituents. The wave functions could be found from the eigenvalue equation

$$H_\lambda |\Psi\rangle = E |\Psi\rangle, \quad (1)$$

where H_λ denotes the QCD Hamiltonian corresponding to a finite renormalization group parameter λ . This parameter limits momentum transfers in effective interactions. Therefore, and if the renormalization group equations for H_λ can be solved to a reasonable accuracy using a series expansion in powers of an effective coupling constant, g_λ , and if λ is brought that way down to scales on the order of hadronic masses, a finite computer code has a chance to find solutions to Eq. (1) and effectively solve the initial theory. Questions concerning the ground state structure, such as symmetry breaking, should in principle be answerable by introducing new Hamiltonian counterterms that remove dependence of the spectrum and other observables on the infinitesimally small cutoff parameter δ^+ that limits the constituents' longitudinal momenta, $k^+ = k^0 + k^3$, from below. Alternatively, the lower bound can also be introduced through factors $r_\delta(x)$ that suppress interactions whenever frac-

tions $x = k^+/P^+$ are smaller than a minuscule dimensionless parameter δ [2], where P^+ is the total momentum carried by particles that a Hamiltonian term annihilates or creates. Each and every interaction may have different P^+ and by using the ratio x one can preserve explicit boost invariance in the renormalization group equations for H_λ . The latter are derived from the relation

$$a_\lambda^\dagger = \mathcal{U}_\lambda a_\infty^\dagger \mathcal{U}_\lambda^\dagger, \quad (2)$$

that changes the original creation (as well as annihilation) operators from the canonical theory, $a_\infty^\dagger$, to the creation operators for effective particles that are denoted by $a_\lambda^\dagger$. The renormalization group parameter λ has interpretation of the inverse size of effective gluons, because all interaction vertices in H_λ contain form factors, f_λ , with momentum width λ .

2. Three-gluon vertex [3]

The third order perturbative results for g_λ are shown in Fig. 1 at the end of this note. The λ -dependent three-particle vertex has the form $W_\lambda + W_\lambda^\dagger$, where $W_\lambda = \sum_{i=1}^4 W_{i\lambda}$, and each and every $W_{i\lambda}$ for $i = 1, \dots, 4$ has the structure

$$W_{i\lambda} = \int [123] \tilde{\delta} f_\lambda V_{i\lambda} Y_{i123} a_{\lambda 1}^\dagger a_{\lambda 2}^\dagger a_{\lambda 3}. \quad (3)$$

$[123]$ denotes the integration measure over the particle three-momenta, $\tilde{\delta}$ is the three-momentum conservation δ -function, $V_{i\lambda} = V_{i\lambda}(x_1, \kappa_{12}^\perp)$ is the effective vertex function of the arguments

*Supported by KBN Grant No. 2P03B01618. E-mail: stglazek@fuw.edu.pl

$x_1 = k_1^\perp/k_3^\perp$ and $\kappa_{12}^\perp = x_2 k_1^\perp - x_1 k_2^\perp$, while f_λ is the vertex form factor from the renormalization group procedure, $f_\lambda = \exp[-\kappa_{12}^{\perp 2}/(x_1 x_2 \lambda^2)]^2$. Y_{i123} for $i = 1, 2, 3$ denote the appropriate color, spin and momentum dependent factors that characterize interactions resulting from the Lagrangian term $gi\partial_\mu A_\nu^a [A^\mu, A^\nu]^a$. Y_{4123} denotes a new structure that emerges from the canonical theory through renormalization group equations but is absent in the bare canonical Lagrangian. Y_{4123} vanishes as $|\kappa_{12}^\perp|^3$ for $\kappa_{12}^\perp \rightarrow 0$ and it does not contribute to the logarithmic running of g_λ . All $V_{i\lambda}(x_1, 0^\perp)$ with $i = 1, 2, 3$ vary equally with λ and the running coupling constant is defined using $W_\lambda(x) \equiv V_{i\lambda}(x, 0^\perp) - V_{i\lambda_0}(x, 0^\perp)$. Namely,

$$W_\lambda(x) = -\frac{g^3}{48\pi^2} N_c [11 + h(x)] \ln \frac{\lambda}{\lambda_0}, \quad (4)$$

describes dependence on λ of all three functions

$$V_{i\lambda}(x, 0^\perp) \equiv g_0 V = g \tilde{r}_\delta(x) + W_\lambda(x) \quad (5)$$

for finite x and λ in the limit $\delta \rightarrow 0$. g_0 is defined by the relation

$$g_0 = \lim_{\delta \rightarrow 0} V_{i\lambda_0}(1/2, 0^\perp). \quad (6)$$

Equation (5) does not display the contribution from quarks. The function $\tilde{r}_\delta(x) = r_\delta(x)r_\delta(1-x)$ depends on the regulating factors that were inserted in the canonical Hamiltonian of QCD to suppress the region of small longitudinal momentum fractions, and $\tilde{r}_\delta(x)$ converges pointwise to 1 when δ is sent to 0.

The Hamiltonian gluon coupling strength V of Eq.(5), depends on x through $h(x)$ and the latter depends on the initial Hamiltonian and small- x regularization factor r_δ . Figure 1 on the next page illustrates results obtained for V as function of x for various values of λ assuming $\alpha_0 = g_0^2/(4\pi) = 0.1$ at $\lambda_0 = 100$ GeV, and $N_c = 3$, in three cases, a), b), and c), corresponding to:

$$r_\delta(x) = x/(x + \delta) \quad , \quad (7)$$

$$r_\delta(x) = \theta(x - \delta) \quad , \quad (8)$$

$$r_\delta(x) = x^\delta \theta(x - \epsilon) \quad , \quad (9)$$

in the limit $\delta \rightarrow 0$. The sharp cutoff case b) is visibly different from c) that produces $h(x) = 0$ for $\epsilon/\delta \ll 1$, but the continuous case a) differs from c) only by 8%.

The case c) with $h(x) = 0$ corresponds to the standard asymptotic freedom result [4] in the sense that it leads to

$$\lambda \frac{d}{d\lambda} g_\lambda = \beta_0 g_\lambda^3 \quad , \quad (10)$$

with

$$\beta_0 = -\frac{11N_c}{48\pi^2} \quad , \quad (11)$$

which is equal to the β -function coefficient in Feynman calculus in QCD. Thus, if one equates the Hamiltonian form factor width λ with the running momentum scale in Feynman diagrams, the standard result from off-shell $\mathcal{S}$ -matrix calculus would be recovered in effective Hamiltonians if $h(x) = 0$, cf. [5,6].

This result explains that asymptotic freedom of effective gluons is a consequence of that a single effective gluon contains a pair of bare gluons. This component amplifies the strength with which effective gluons split into effective gluons when λ gets smaller. The fact that such running coupling constant appears in the renormalized Hamiltonians with small cutoffs, explains also why it is possible that a running coupling in the scattering processes provides a good approximation to physics. Namely, in perturbative description of processes characterized by a physical momentum scale Q , using $H_\lambda(a_\lambda)$, there will appear powers of g_λ and $\ln Q/\lambda$. For $Q/\lambda = 1$, $\ln Q/\lambda = 0$, so that the theoretical predictions will have a power expansion in the asymptotically free running coupling constant g_Q .


3. Conclusion

One can trace the origin of why the factor $r_\delta \sim x^\delta$ in LF QCD Hamiltonians provides a connection with the covariant Lagrangian calculus to the success of dimensional regularization in evaluating Feynman diagrams: the parameter δ is analogous to the deviation of the number of dimensions from 4. Although the ultra-violet Hamiltonian counterterms in canonical LF

QCD are much more complicated [2] than in the Lagrangian calculus, $r_\delta \sim x^\delta$ avoids the anomalous $h(x)$ in the effective gluon vertex in small- λ Hamiltonians once the latter are already made insensitive to the ultraviolet regularization via renormalization group. While $h(x)$ may appear undesirable in Feynman calculus, Fig. 1 shows that for r_δ from Eqs. (7) and (8), $h(x)$ may lead to potentially useful dynamical suppression of extreme values of x in gluon interactions and help in building a selfconsistent approximation scheme for solving Eq. (1) with small values of λ .

The general problem of deriving covariant results for observables in the LF Hamiltonian calculus with small λ , cannot be solved by showing a connection with Feynman diagrams that are limited to perturbation theory, when one wants to solve Eq. (1) non-perturbatively. Fortunately, the same renormalization group procedure for effective particles is capable, at least in principle, of providing solutions for all generators of the Poincaré algebra [7] and enabling us to study the symmetry beyond perturbation theory.

Fig. 1 Variation of the effective gluon vertex strength V from Eq. (5) with the width λ and gluon momentum fraction x . λ changes from $\lambda_0 = 100$ GeV down to 100 MeV, for three different small- x regularizations; a) Eq. (7), b) Eq. (8), and c) Eq. (9). Part d) shows dependence of $V(1/2)$ on λ for the three cases, correspondingly. Note the dynamical suppression of the effective gluon coupling strength for extreme values of x in cases a) and b). The case c) matches the standard Lagrangian running coupling constant result obtained from Feynman diagrams.



REFERENCES

1. K. G. Wilson et al., Phys. Rev. D**49** (1994) 6720.
2. See e.g. St. D. Glazek, Nucl. Phys. B (Proc. Suppl.) **90** (2000) 175.
3. St. D. Glazek, Phys. Rev. D**63** (2001) 116006.
4. H. D. Politzer, Phys. Rev. Lett. **30** (1973) 1346; D. J. Gross and F. Wilczek, Phys. Rev. Lett. **30** (1973) 1343.
5. C. B. Thorn, Phys. Rev. D**20** (1979) 1934.
6. R. J. Perry, Phys. Lett. B**300** (1993) 8.
7. St. D. Glazek, T. Masłowski, hep-th/0110185.